

**Fire history of a western Montana
ponderosa pine grassland:
A pilot study**

Don V. Gayton¹, Marc H. Weber², Mick Harrington³, Emily K. Heyerdahl⁴, Elaine K. Sutherland⁵, Bob Brett⁶, Cindy Hall⁷, Michael Hartman⁸, Liesl Peterson⁹, Carolynne Merrel¹⁰

¹Forest Research Extension Partnership, Nelson, B.C. ²Fire Sciences Lab, Missoula, MT ³Fire Sciences Lab ⁴Fire Sciences Lab ⁵Forestry Sciences Lab, Missoula, MT ⁶Snowline Ecological Consulting, Whistler, BC ⁷Natural Elements Consulting, Rossland, BC ⁸World Data Center for Paleoclimatology, Boulder, CO ⁹Colorado College, Colorado Springs, CO ¹⁰University of Idaho, Moscow, ID

Introduction

A primary goal in the management of forests and grasslands is to maintain community structure and disturbance processes within their historical range of variation. If, within a managed ecosystem, either is found to lie outside that range, restoration may be necessary. Both maintenance and restoration are currently guided by the principles of ecosystem management, which relies on knowledge of both historical processes and current ecosystem conditions (Forest Ecosystem Management Team 1993). In ecosystems historically sustained by fire, site-specific fire regime data can be combined with information on present composition and structure to design ecologically appropriate restoration and management prescriptions. While this approach to restoring fire-adapted ecosystems is appropriate for many publicly managed forests, it is actually mandated for U.S. Forest Service-designated Research Natural Areas (RNA). Research Natural Areas

are established as examples of forests or grasslands that most closely represent historical vegetation and wildlife habitat and that are largely products of natural disturbance processes and ecosystem succession (U.S. Department of Agriculture, 1994).

The 110 ha Sawmill Creek Research Natural Area is 16 km southeast of Stevensville, Montana, in the Bitterroot National Forest. This area was designated a RNA in 1992 to preserve, through natural disturbance processes or their surrogates, its native bunchgrass communities and the old growth ponderosa pine (*Pinus ponderosa*) /Douglas-fir (*Pseudotsuga menziesii*) stands that grow in several narrow, north-south ravines (Figure 1). However, the vegetation composition and structure of the RNA has changed in recent decades, and now may be in a condition outside its historical range of variation. Specifically, numerous Douglas-fir trees have established in the understory of the ravine forests, and Rocky Mountain juniper has encroached on the grasslands. We speculate fire exclusion was a likely cause of the recent changes in structure and composition, as it has been elsewhere in the interior west (e.g., Weaver 1959, 1961, Cooper 1960, Covington and Moore 1994). Surface fires were historically frequent in ponderosa pine forests near the RNA (Barrett 1981), and some of the large ponderosa pine trees in the ravines at Sawmill have multiple fire scars, suggesting that the RNA historically sustained frequent surface fires as well. Forest Service Fire Reports initiated in the early 1900's contain no record of significant wildfire activity in the RNA, although a 1923 fire burned 300 acres near the northeast boundary (Johnson and Stewart 1998), and could have burned through the RNA's upper grasslands.

Our objective for this pilot study was to reconstruct a preliminary, multi-century history of surface fires at the Sawmill Creek RNA from fire scars on ponderosa pine trees. This fire history will provide evidence of the frequency of past fires that sustained the forest and grassland communities in the RNA, which may be used to help guide future management of the RNA.



Figure 1. Upper Sawmill Creek RNA, showing forested ravines and recent invasion of grasslands.

Study area

Elevation at the Sawmill Creek RNA ranges from 1400 to 1650 m. The slopes are primarily southerly, but are bisected by moderately steep, north-south ravines. The climate is generally continental, with hot, dry summers and cold winters (Figure 2). Monthly precipitation peaks in May-June, followed by temperature peaks in July and August.

The RNA has five distinct plant communities (Sikkink 1997):

- Grassland (bluebunch wheatgrass, Idaho fescue, rough fescue, arrowleaf balsamroot).
- Shrub Savanna (big sagebrush, arrowleaf balsamroot, rough fescue),
- Mixed Tree-Shrub Savanna (ponderosa pine, Douglas-fir, blue-

bunch wheatgrass, snowberry, arrowleaf balsamroot).

- Canyon Forest (ponderosa pine, Douglas-fir, Rocky Mountain juniper, pinegrass, Idaho fescue,
- Riparian forest (ponderosa pine, Douglas-fir, Rocky Mountain maple, mockorange, gray alder).




Figure 2. Climate at Stevensville, Montana, 16 km northwest of Sawmill Creek RNA (1911-2002, Western Regional Climate Center 2003). Bars indicate total monthly precipitation; lines indicate maximum, mean, and minimum monthly temperatures.

The Grassland community occurs on the upper slopes and drier aspects of the RNA: Shrub Savanna and Mixed Tree-Shrub Savanna are found on the midslopes, and Canyon Forest on the lower slopes and mesic aspects of the ravines. Riparian Forest communities form narrow ribbons along the bottoms of the ravines. We sampled in the Tree-Shrub Savanna and Canyon Forest types.

The RNA was influenced by twentieth-century human activities before its designation in 1992. Prior to 1945, the RNA was subject to unregulated livestock grazing, followed by permitted grazing until 1961 (Johnson and Stewart 1998). In the 1950's, standing snags were cut to reduce the occurrence of lightning fires. In the 1960's,

ring-boundary (dormant season) scars (12 and 9 scars, respectively).

Discussion

The frequency of past surface fires at the RNA is similar to that reconstructed in similar forest types elsewhere in this region (e.g., Barrett and Arno 1982, Arno *et al.* 1997). For ~200 years prior to 1900, surface fires burned at Sawmill Creek RNA every 13 years, on average, with the longest interval being 32 years. However, similar to other sites in the region, surface fires in our study area ceased abruptly prior to 1900. The last fire reconstructed at the RNA occurred 114 years ago, more than triple the longest previous interval. Significant changes in plant community composition and structure would be expected as a result of this extended fire-free period.




Figure 5. Composite fire-return intervals at Sawmill Creek RNA. The box encloses the 25th to 75th percentiles, while the whiskers enclose the 10th to 90th percentiles of the distribution of fire return intervals. The vertical line indicates the median, and values falling outside the 10th to 90th percentiles are shown as circles.

Sikkink's (1997) comprehensive analysis of the Sawmill RNA vegetation indicates that currently over twice as many Douglas-fir occupy the site than the more fire-adapted ponderosa pine. Ponderosa pine's fire tolerance advan-

tage over Douglas-fir, especially in the smaller size classes (Fischer and Bradley 1987; Kalabokidis and Wakimoto 1992) was likely a primary factor in the development of a ponderosa pine-dominated stand during the period of regular fire occurrence. Old ponderosa pine still dominates the overstory (i.e., trees >50 cm), but Douglas-fir, overwhelmingly, dominates all younger, smaller size classes (Figure 6). In addition, Douglas-fir, with a high percentage of dwarf-mistletoe infection, appears to be more disease-prone than ponderosa pine at this site (data on file, M.G. Harrington, Fire Sciences Lab, Missoula MT).




Figure 6. Density by size class of Douglas-fir and ponderosa pine in the upland forest communities at Sawmill Creek RNA (adapted from Sikkink, 1997). Note difference in Y axis scales.

The recent increase in forest density, and the shift toward less fire-tolerant species is consistent with our interpretation of a previously fire-suppressed, open forest ecosystem now experiencing forest ingrowth. Without a return to the short fire intervals that controlled stand density and reduced fuels, existing old-growth trees will likely be lost in a future, stand-replacing wild-fire or suffer stress due to competition for limited site resources. Addition-

ally, the current condition will likely continue to diminish the regeneration potential of ponderosa pine and the vigor of associated shade-intolerant understory species.

The re-introduction of fire and/or fire surrogates to the Sawmill Creek RNA appears to be an appropriate strategy for restoring this area to its historical condition. However, simply because a forest or grassland was maintained historically by fire does not dictate the immediate use of fire as a restoration tool. For ecosystems adapted to low-severity, surface fires, where an accepted departure from the historical or desired ecological conditions are relatively small, prescribed fires can be a viable replacement for natural fires. Prescribed fires can be effective in reducing the density of juvenile, fire-sensitive trees and in consuming excess organic matter. Conversely, if the current departure is large, as at the Sawmill RNA, returning fire alone may result in undesired consequences (e.g., Tiedeman *et al* 2000). Attempting to reduce excess juvenile and pole-sized trees and substantial amounts of organic debris with prescribed fire can lead to mortality of the favored old growth cohort. Preliminary experiments with mechanical tree and fuel

reduction have proven successful in reducing this risk, even though some regard it as less ecologically acceptable than the use of fire treatments alone (Fiedler *et al* 1996). Equipped with information about the historical fire regime, the manager can develop an appropriate plan that replicates the historical disturbance process that once sustained these forest and grassland ecosystems. The manager might also choose to more closely approximate the range of historical variation by varying both the fire return interval as well as the season of burning.

Since the completion of the pilot study, we have obtained fire-scarred sections from an additional eighteen ponderosa pine trees at the RNA. We will cross-date these sections to yield a more complete reconstruction of fire history that will be used in the current effort to restore this RNA to its historical ecosystem structure and composition.

Acknowledgments

We thank Jim Speer for organizing the 13th North American Dendroecological Field Week, during which this study was conducted, and the USDA Forest Service Rocky Mountain Research Station, for logistical support.

Literature Cited

- Arno, S.F. 19980. Forest fire history of the Northern Rockies. *Journal of Forestry*. 78(8):460-465.
- Arno, S.F. 1976. The historical role of fire on the Bitterroot National Forest. USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah. Res. Pap. INT-187 pp. 1-29.
- Arno, S.F., *et al*. 1997. Old growth ponderosa pine and western larch stand structures: Influence of pre-1900 fires and fire exclusion. Research Paper INT-RP-495, USDA, Intermountain Research Station Ogden, UT. 20 pps.
- Barrett, S.W. and Arno, S.F. 1982. Indian fires as an ecological influence in the Northern Rockies. *J. Forestry* 80:647-665.

- Cook, E. R., *et al.* 1999. Drought reconstructions for the continental United States. *Journal of Climate*. 12:1145-1162. Data archived at the World Data Center for Paleoclimatology, Boulder, Colo..
- Cooper, C.F. 1960. Changes in vegetation, structure, and growth of southwestern pine forests since white settlement. *Ecological Monographs*. 30:129-164.
- Covington, W.W. and Moore, M.M. 1994. Postsettlement changes in natural fire regimes and forest structure: Ecological restoration of old-growth ponderosa pine forest. In: Sampson, R.N. and Adams, D.L., eds. *Assessing the forest ecosystem health in the Inland West*. *Journal of Sustainable Forestry* 2:153-181.
- Dieterich, J.H. 1980. The composite fire interval: A tool for more accurate interpretation of fire history. Pages 8-14 in M.A. Stokes and J.H. Dieterich, technical coordinators. *Proceedings of the Fire History Workshop, 20-24 Oct. 1980, Tucson*. General Technical Report RM-81. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo.
- Fieldler, C.E., *et al.* 1996. Flexible silviculture and prescribed burning approaches for improving health of ponderosa pine forests. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, Colo. Gen. Tech. Rep. RM-GTR-278 pp 69-74.
- Fischer, W. C., and Bradley, A. F. 1987. Fire ecology of western Montana forest habitat types. Gen. Tech. Rep. INT-223. USDA Forest service. Intermountain Research Station, Ogden, Utah. 94 p.
- Forest Ecosystem Management Assessment Team. 1993. *Forest ecosystem management: An ecological, economic, and social assessment*. Washington, D.C.:USDA, Forest Service; US Department of Commerce, National Oceanic and Atmospheric Administration, US Department of Interior. Bureau of Land Management. U.S. Fish and Wildlife Service, and National Park Service; and Environmental Protection Agency.
- Grissino-Mayer, H.D. 1995. Tree-ring reconstructions of climate and fire history at El Malpais National Monument, New Mexico. Ph.D. dissertation. Tucson: University of Arizona. 407 p.
- Habeck, J.R. et al. 1992. USDA Forest Service establishment record for Sawmill Creek Research Natural Area within Bitterroot National Forest, Ravalli County, Montana, unpublished, 20 p.
- Heyerdahl, E.H. *et al.* 2002. Spatial controls of historical fire regimes: A multiscale example from the Interior West, USA. *Ecology* 82:660-678.
- Holmes, R.L. 1983. Computer-assisted quality control in tree-ring dating and measurement. *Tree-Ring Bulletin*. 43:69-78.
- Johnson, J.L. and Stewart, C. 1998. Sawmill Research Natural Area Management Plan (Draft). Bitterroot National Forest, 29 pp.
- Kalabokidis, K. D., and Wakimoto R. H. 1992. Prescribed fire in uneven-aged stand management of ponderosa pine/Douglas-fir forests. *Journal of Environmental Management* 43 (3): 221-235.
- Sikkink, P.G. 1997. Ecological characterization and restoration alternatives for Sawmill Creek Research Natural Area, Bitterroot National Forest, Montana. Missoula, MT: University of Montana, Final Report INT-95076-RJVA. 108 pp.
- Tiedemann, A. *et al.* 2000. Solution of Forest Health Problems with Prescribed Fire: Are Forest Productivity and Wildlife at Risk? *Forest Ecology and Management* 127 (2000):1-18.

- USDA, Forest Service. 1994. Forest Service Manual, reference 4063. Research Natural Areas. WO Amendment 4000-94-2, effective May 4, 1994. Washington, DC:USDA, Forest Service.25 p.
- Weaver, H. 1959. Ecological changes in the ponderosa pine forest of the Warm Springs Indian Reservation in Oregon. *Journal of Forestry*. 57:15-20.
- Weaver, H. 1961. Ecological changes in the ponderosa pine forest of Cedar Valley in southern Washington. *Ecology*. 42:416-420.
- Western Regional Climate Center, ed. 2003. Montana Climate Summaries [Homepage of Western Regional Climate Center], [Online]. URL: <http://www.wrcc.dri.edu/summary/climsmt.html>